

LIST OF REFERENCES

- Alhonen, P., and Haavisto, M-L. (1969). The biostratigraphical history of Lake Otalampi in southern Finland, with special reference to the remains of subfossil midge fauna. Bulletin of the Geological Society of Finland 41: 157-164.
- Allee, W.C., Emerson, A.E., Park, O., Park, T., and Schmidt, K.P. (1949). Principles of Animal Ecology. W.B. Saunders, London. 837pp.
- Andersen, F.S. (1937). Ueber die Metamorphose der Ceratopogoniden und Chironomiden Nordost-Grønlands. Meddelelser om Grønland 116(1): 1-95.
- Andersen, F.S. (1938). Spätglaziale Chironomiden. Meddelelser Dansk geologisk Forening 9: 320-326.
- Andersen, F.S. (1943). *Dryadotanytarsus edentulus* n.gen. et sp. (Dipt., Chiron.) from the late glacial period in Denmark. Entomologiske Meddelelser 23: 174-178.
- Andersen, F.S. (1946). East Greenland lakes as habitats for chironomid larvae. Studies on the systematics and biology of Chironomidae. II. Meddelelser om Grønland 100: 1-69.
- Andrews, D., and Rigler, F.H. (1985). The effects of an arctic winter on benthic invertebrates in the littoral zone of Char Lake, Northwest Territories. Canadian Journal of Zoology 63: 2825-2834.
- Ankar, S., Andersin, A-B., Lassig, J., Norling, L., and Sandler, H. (1979). Methods of studying benthic macrofauna, an intercalibration between two laboratories in the Baltic Sea. Finnish Marine Research 246: 147-160.
- Bacon, C.R. (1983). Eruptive history of Mount Mazama and Crater Lake Caldera, Cascade Range, U.S.A. Journal of Volcanology and Geothermal Research 18: 57-115.
- Banner, A., Pojar, J., and Trowbridge, R. (1983). Ecosystem classification of the Coastal Western Hemlock Zone, Queen Charlotte Island subzone (CWHg), Prince Rupert Forest Region, British Columbia. (author's preliminary draft). Ministry of Forests, Province of British Columbia
- Barnosky, C.W. (1984). Late Pleistocene and early Holocene environmental history of southwestern Washington State, U.S.A. Canadian Journal of Earth Sciences 21: 619-629.
- Barton, D.R., and Smith, S.M. (1984). Insects of extremely small and extremely large aquatic habitats. pp. 456-483, In: The Ecology of Aquatic Insects. (V.H. Resh and D.M. Rosenberg, Eds.), Praeger Publ., New York.
- Bass, D. (1986). Habitat ecology of chironomid larvae of the Big Thicket streams. Hydrobiologia 134: 29-41.
- Baust, J.G., and Edwards, J.S. (1979). Mechanisms of freezing tolerance in an Antarctic midge, *Belgica antarctica*. Physiological Entomology 4: 1-5.

- Beck, W.M., Jr. (1980). Interesting new chironomid records for the southern United States (Diptera: Chironomidae). Journal of the Georgia Entomological Society 15: 69-73.
- Bennike, O. (personal communication). Geological Museum, Copenhagen University, Øster Voldegade 5-7, DK-1350 Copenhagen, Denmark.
- Berglund, B.E., and Digerfeldt, G. (1970). A palaeoecological study of the Late-Glacial lake at Torreberga, Scania, south Sweden. Oikos 21: 98-128.
- Bliss, L.C. (Ed.). (1977). Truelove Lowland, Devon Island, Canada: A high arctic ecosystem. University of Alberta Press, Edmonton.
- Bode, R.W. (1983). Larvae of North American *Eukiefferiella* and *Tvetenia* (Diptera: Chironomidae). New York State Museum Bulletin 452: 1-40.
- Borkent, A. (1979). Systematics and bionomics of the species of the subgenus *Schadonophasma* Dyar and Shannon (*Chaoborus*, Chaoboridae, Diptera). Quaestiones Entomologicae 15: 122-255.
- Borkent, A. (1984). The systematics and phylogeny of the *Stenochironomus* complex (*Xestochironomus*, *Harrisius*, and *Stenochironomus*) (Diptera: Chironomidae). Memoirs of the Entomological Society of Canada 128: 1-269.
- Brehm, V., Krasske, G., and Krieger, W. (1948). Subfossile tierische Reste und Algen im Schwarzsee bei Kitzbühel. Osterreichische botanische Zeitschrift 95: 74-83.
- Brewer, M.C. (1958). The thermal regime of an arctic lake. Transactions of the American Geophysical Union 39: 278-284.
- Brinkhurst, R.O. (1974). The benthos of lakes. MacMillan Press, London.
- Broccoli, A.J., and Manabe, S. (1987). The effects of the Laurentide Ice Sheet on North American climate during the last glacial maximum. Géographie physique et Quaternaire 41: 291-299.
- Brodin, Y. (1986). The postglacial history of Lake Flarken, southern Sweden, interpreted from subfossil insect remains. Internationale Revue der gesamten Hydrobiologie 71: 371-432.
- Brooke, R.C., Peterson, E.B., and Krajina, V.J. (1970). The subalpine mountain hemlock zone. Ecology of Western North America 2: 147-349.
- Brundin, L. (1948). Über die Metamorphose der Sectio Tanytarsariae connectentes (Dipt., Chironomidae). Arkiv for Zoologi 41A: 1-22.
- Brundin, L. (1949). Chironomiden und andere Bodentiere der südschwedischen Urgebirgsseen. Report of the Institute of Freshwater Research, Drottningholm 30: 1-914.
- Brundin, L. (1951). The relation of O₂-microstratification at the mud surface to the ecology of the profundal bottom fauna. Report of the Institute of Freshwater Research, Drottningholm 32: 32-42.

- Brundin, L. (1956). Die bodenfaunistischen Seetypen und ihre Anwendbarkeit auf die Südhalbkugel. Zugleich eine Theorie der produktionsbiologischen Bedeutung der glazialen Erosion. Report of the Institute of Freshwater Research, Drottningholm 37: 186-235.
- Brundin, L. (1958). The bottom faunistic lake type system and its application to the southern hemisphere. Moreover a theory of glacial erosion as a factor of productivity in lakes and oceans. Verhandlungen der internationalen Vereinigung theoretische und angewandte Limnologie 13: 288-297.
- Brundin, L. (1983). The larvae of Podonominae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. Entomologica scandinavica Supplement 19: 23-31.
- Bryce, D. (1962). Chironomidae (Diptera) from fresh water sediments with special reference to Malham Tarn (Yorks). Transactions of the Society for British Entomology 15: 41-54.
- Brylinsky, M., and Mann, K.H. (1973). An analysis of factors governing productivity in lakes and reservoirs. Limnology and Oceanography 18: 1-14.
- Bryson, R.A. (1966). Air masses, streamlines, and the boreal forest. Geographical Bulletin 8: 228-269.
- Bunnell, F.L., McNay, R.S., and Shank, C.C. (1985). Trees and snow: the deposition of snow on the ground - a review and quantitative synthesis. Research, Ministries of Environment and Forests. IWIFR-17, Victoria, B.C.
- Butler, M., Miller, M.C., and Mozley, S. (1981). Macroinvertebrates. pp. 297-339, In: Limnology of Tundra Ponds, Barrow, Alaska. (J.E. Hobbie, Ed.), US/IBP Synthesis Series 13. Dowden, Hutchinson and Ross, Stroudsburg, Pennsylvania.
- Calder, J.A., and Taylor, R.L. (1968). Flora of the Queen Charlotte Islands. Part 1. Canada Department of Agriculture Monograph 4: 1-659.
- Cannings, R.A. (1975a). Some Chironomidae (Diptera) new to British Columbia and Canada. Journal of the Entomological Society of British Columbia 72: 23-25.
- Cannings, R.A. (1975b). A new species of *Chironomus* (Diptera: Chironomidae) from saline lakes in British Columbia. Canadian Entomologist 107: 447-450.
- Cannings, R.A., and Scudder, G.G.E. (1978). The littoral Chironomidae (Diptera) of saline lakes in central British Columbia. Canadian Journal of Zoology 56: 1144-1155.
- Chang, P.S. (1975). Emergence of insects from Shell Lake and Explosive Lake, MacKenzie Delta, N.W.T. Canada Fisheries and Marine Service Resource Development Technical Report 554: 1-14.
- Clague, J.J. (1981). Late Quaternary geology and geochronology of British Columbia. Part 2. Summary and discussion of radiocarbon-dated Quaternary history. Geological Survey of Canada Paper 80-35.

- Clague, J.J., Mathewes, R.W., and Warner, B.G. (1982). Late Quaternary geology of eastern Graham Island, Queen Charlotte Islands, British Columbia. Canadian Journal of Earth Sciences 19: 1786-1795.
- Coffman, W.P., Cranston, P.S., Oliver, D.R., and Sæther, O.A. (1986). The pupae of Orthoclaadiinae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. Entomologica scandinavica Supplement 28: 147-296.
- Coffman, W.P., and Ferrington, L.C., Jr. (1984). Chironomidae. pp. 551-652, In: An Introduction to the Aquatic Insects of North America. 2nd ed. (R.W. Merritt and K.W. Cummins, Ed.), Kendall/Hunt Publishing, Dubuque, Iowa.
- Corbet, P.S. (1972). The microclimate of arctic plants and animals, on land and in freshwater. Acta arctica 18: 1-43.
- Cranston, P.S. (1982a). The metamorphosis of *Symposiocladius lignicola* (Kieffer) n.gen., n.comb., a wood-mining Chironomidae (Diptera). Entomologica scandinavica 13: 419-429.
- Cranston, P.S. (1982b). A key to the larvae of the British Orthoclaadiinae (Chironomidae). Freshwater Biological Association, Scientific Publication 45: 1-152.
- Cranston, P.S., and Oliver, D.R. (in press). Aquatic xylophagous Orthoclaadiinae - systematics and ecology. Spixiana
- Cranston, P.S., Oliver, D.R., and Sæther, O.A. (1983). The larvae of Orthoclaadiinae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. Entomologica scandinavica Supplement 19: 149-291.
- Crisman, T.L. (1978). Reconstruction of past lacustrine environments based on the remains of aquatic invertebrates. pp. 69-101, In: Biology and Quaternary environments. (D. Walker and J.C. Guppy, Eds.), Australian Academy of Science, Canberra City.
- Damman, A.W.H. (1965). The distribution patterns of northern and southern elements in the flora of Newfoundland. Rhodora 67: 363-392.
- Danks, H.V. (1971a). Overwintering of some north temperate and arctic Chironomidae. II. Chironomid biology. Canadian Entomologist 103: 1875-1910.
- Danks, H.V. (1971b). Spring and early summer temperatures in a shallow arctic pond. Arctic 24: 113-123.
- Danks, H.V. (1971c). Overwintering of some north temperate and arctic Chironomidae. I. The winter environment. Canadian Entomologist 103: 589-604.
- Danks, H.V. (1981). Arctic Arthropods: A review of systematics and ecology with particular reference to the North American fauna. Entomological Society of Canada, Ottawa. 608p.
- Danks, H.V., and Oliver, D.R. (1972a). Seasonal emergence of some high arctic Chironomidae (Diptera). Canadian Entomologist 104: 661-686.

- Danks, H.V., and Oliver, D.R. (1972b). Diel periodicities of emergence of some high arctic Chironomidae (Diptera). Canadian Entomologist 104: 903-916.
- Davies, I.J. (1975). Selective feeding in some arctic Chironomidae. Verhandlungen der Internationalen Vereinigung für theoretische und angewandte Limnologie 19: 3149-3154.
- Davis, M.B. (1984). Climatic instability, time lags, and community disequilibrium. pp. 269-284, In: Community Ecology. (J. Diamond and E.J. Case, Eds.), Harper and Row, N.Y.
- Davis, M.B., Moeller, R.E., and Ford, J. (1984). Sediment focusing and pollen influx. pp. 261-293, In: Lake sediments and environmental history. (E.Y. Haworth and J.W.G. Lund, Eds.), Leicester University Press, U.K.
- Davis, R.B., and Anderson, D.S. (1985). Methods of pH calibration of sedimentary diatom remains for reconstructing history of pH in lakes. Hydrobiologia 120: 69-87.
- de March, L., de March, B., and Eddy, W. (1978). Limnological, fisheries, and stream zoobenthic studies at Stanwell-Fletcher Lake, a large high arctic lake. Arctic Islands Pipelines Program, Preliminary Report 1977. Department of Indian and Northern Affairs Publication No. QS-8160-004-EE-A1, Ottawa.
- Deevey, E.S., Jr. (1942). Studies on Connecticut lake sediments. III. The biostratonomy of Linsley Pond. American Journal of Science 240: 233-264, 313-324.
- Deevey, E.S., Jr. (1955a). The obliteration of the hypolimnion. Memorie dell'Istituto italiano di Idrobiologia, Supplement 8: 9-38.
- Deevey, E.S., Jr. (1955b). Paleolimnology of the Upper Swamp deposit, Pyramid Valley. Records of the Canterbury Museum 6: 291-344.
- Deevey, E.S., Jr. (1969). Coaxing history to conduct experiments. BioScience 19: 40-43.
- Donald, D.B. (personal communication). Water Planning and Management Branch, Western and Northern Region, Inland Waters Directorate, Environment Canada, 1901 Victoria Avenue, Regina, Saskatchewan, Canada S4P 3R4.
- Downes, J.A. (1962). What is an arctic insect? Canadian Entomologist 94: 143-162.
- Downes, J.A. (1964). Arctic insects and their environment. Canadian Entomologist 96: 279-307.
- Dunbar, M. (1985). Sea ice and climatic change in the Canadian arctic since 1800. Syllogeus 55: 107-119.
- Driver, E.A. (1977). Chironomid communities in small prairie ponds: some characteristics and controls. Freshwater Biology 7: 121-133.
- Efford, I.E. (1967). Temporal and spatial differences in phytoplankton productivity in Marion Lake, British Columbia. Journal of the Fisheries Research Board of Canada 24: 2283-2307.

- Efford, I.E., and Hall, K.J. (1975). Marion Lake - An analysis of a lake ecosystem. pp. 199-219. In: Energy flow, its biological dimensions. A summary of the IBP in Canada, 1964-1974. (T.W.M. Cameron and L.W. Billingsley, Eds.), Royal Society of Canada, Ottawa.
- Ekman, S. (1915). Die Bodenfauna des Vättern, qualitativ und quantitativ untersucht. Internationale revue der gesamten Hydrobiologie und Hydrographie 7: 146-204, 275-425.
- Elgmork, K., and Sæther, O.A. (1970). Distributions of invertebrates in a high mountain brook in the Colorado Rocky Mountains. University of Colorado Studies Series in Biology 31: 1-55.
- Environment Canada. (1982). Canadian climate normals (1951-1980). Vol. 2, 3, 4, and 6. Atmospheric Environment Service, Ottawa, Canada.
- Farley, A.L. (1979). Atlas of British Columbia - People, environment and resource use. University of British Columbia Press, Vancouver, 136pp.
- Fitkau, E.J., and Reiss, F. (1978). Chironomidae. pp. 404-440. In: Limnofauna europaea, 2nd ed. (J. Illies, Ed.), Gustav Fischer Verlag, Stuttgart.
- Fittkau, E.J., and Roback, S.S. (1983). The larvae of Tanypodinae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. Entomologica scandinavica Supplement 19: 33-110.
- Fjellberg, A. (1972). Present and late Weichselian occurrence of *Corynocera ambigua* Zett. (Dipt., Chironomidae) in Norway. Norsk Entomologisk Tidsskrift 19: 59-61.
- Flannagan, J.F. (1973). Sorting benthos using floatation media. Fisheries Research Board of Canada Technical Report 354: 1-14.
- Frey, D.G. (1955). Långsee: a history of meromixis. Memorie dell'Istituto italiano di Idrobiologia, Supplement 8: 141-164.
- Frey, D.G. (1964). Remains of animals in Quaternary lake and bog sediments and their interpretation. Ergebnisse der Limnologie 2: 1-114.
- Frey, D.G. (1976). Interpretations of Quaternary paleoecology from Cladocera and midges, and prognosis regarding usability of other organisms. Canadian Journal of Zoology 54: 2208-2226.
- Gams, H., (1927). Die Geschichte der Lunzer Seen, Moore und Wälder. Internationale revue der gesamten Hydrobiologie und Hydrographie 18: 305-387.
- Gannon, J.E. (1971). Two counting cells for the enumeration of zooplankton micro-crustacea. Transactions of the American Microscopical Society 90: 486-490.
- Garrett, C.B.D. (1925). Seventy new Diptera. Cranbrook, B.C. 16 p.
- Goulden, C.E. (1964). Progressive changes in the cladoceran and midge fauna during the ontogeny of Esthwaite Water. Verhandlungen der Internationalen Vereinigung für theoretische und angewandte Limnologie 15: 1000-1005.

- Green, D.G. (1987). Pollen evidence for the postglacial origins of Nova Scotia's forests. Canadian Journal of Botany 65: 1163-1179.
- Gross, H. (1937). Nachweis der Allerödschwankung im süd- und ostbaltischen Gebiet. Botanisches Zentralblatt, Beihandlung, Abteilung B 57: 167-318.
- Günther, J. (1983). Development of Grossensee (Holstein, Germany): variations in trophic status from the analysis of subfossil microfauna. Hydrobiologia 103: 231-234.
- Hågvar, S., and Østbye, E. (1973). Notes on some winter-active Chironomidae. Norsk Entomologisk Tidsskrift 20: 253-257.
- Hall, K.J., and Hyatt, K.D. (1974). Marion Lake (IBP) - from bacteria to fish. Journal of the Fisheries Research Board of Canada 31: 893-911.
- Hamilton, A.L. (1965). An analysis of a freshwater benthic community with special reference to the Chironomidae, 2 parts. Ph.D. thesis, University of British Columbia, Vancouver, Canada.
- Hann, B.J., and Warner, B.G. (1987). Late Quaternary Cladocera from coastal British Columbia, Canada: a record of climatic or limnologic change? Archiv für Hydrobiologie 110: 161-177.
- Hansen, D.C., and Cook, E.F. (1976). The systematics and morphology of the Nearctic species of *Diamesa* Meigen, 1835 (Diptera: Chironomidae). Memoirs of the American Entomological Society 30: 1-203.
- Hansen, K. (1962). The dystrophic lake type. Hydrobiologia 19: 183-191.
- Hare, R.L. (1976). The macroscopic zoobenthos of Parry Sound, Georgian Bay. M.Sc. thesis, University of Waterloo, Waterloo, Canada.
- Hebda, R.J. (1983). Late-glacial and postglacial vegetation history at Bear Cove Bog, northeast Vancouver Island, British Columbia. Canadian Journal of Botany 61: 3172-3192.
- Hebda, R.J., and Mathewes, R.W. (1984). Holocene history of cedar and native indian cultures of the North American Pacific coast. Science 225: 711-713.
- Heikkinen, O. (1984a). Dendrochronological evidence of variations of Coleman Glacier, Mount Baker, Washington, U.S.A. Arctic and Alpine Research 16: 53-64.
- Heikkinen, O. (1984b). Forest expansion in the subalpine zone during the past hundred years, Mount Baker, Washington, U.S.A. Erdkunde 38: 194-202.
- Heikkinen, O. (1984c). Climatic changes during recent centuries as indicated by dendrochronological studies, Mount Baker, Washington, U.S.A. pp. 353-361, In: Climatic changes on a yearly to a millennial basis. (N.-A. Mörner and W. Karlén, Eds.), D. Reidel Publishing.
- Heikkinen, O. (1985). Relationships between tree growth and climate in the subalpine Cascade Range of Washington, U.S.A. Annales Botanici Fennici 22: 1-14.

- Henrikson, L., Olofsson, J.B., and Oscarson, H.G. (1982). The impact of acidification on Chironomidae (Diptera) as indicated by subfossil stratification. Hydrobiologia 86: 223-229.
- Hershey, A.E. (1985a). Littoral chironomid communities in an arctic Alaskan lake. Holarctic Ecology 8: 39-48.
- Hershey, A.E. (1985b). Effects of predatory sculpin on the chironomid communities in an arctic lake. Ecology 66: 1131-1138.
- Heusser, C.J. (1972). Palynology and phytogeographical significance of a late-Pleistocene refugium near Kalaloch, Washington. Quaternary Research 2: 189-201.
- Heusser, L.E. (1983). Palynology and paleoecology of postglacial sediments in an anoxic basin, Saanich Inlet, British Columbia. Canadian Journal of Earth Sciences 20: 873-885.
- Hilsenhoff, W.L. (1966). The biology of *Chironomus plumosus* (Diptera: Chironomidae) in Lake Winnebago, Wisconsin. Annals of the Entomological Society of America 59: 465-473.
- Hilsenhoff, W.L. (1967). Ecology and population dynamics of *Chironomus plumosus* (Diptera: Chironomidae) in Lake Winnebago, Wisconsin. Annals of the Entomological Society of America 60: 1183-1194.
- Hirvenoja, M. (1960). Massenaufreten von *Corynocera ambigua* Zett. (Dipt. Chironomidae) im See Sompiojärvi, Finnish-Lapland. Annales Entomologici Fennici 26: 157-163.
- Hirvenoja, M. (1961). Description of the larva of *Corynocera ambigua* Zett. (Dipt. Chironomidae) and its relation to the subfossil species *Dryadotanytarsus edentulus* Anders. and *D. duffi* Deevey. Annales Entomologici Fennici 27: 105-110.
- Hobbie, J.E., (1973). Arctic limnology: a review. pp. 127-168, In: Alaskan Arctic Tundra. (W.W. Gunn, Ed.) Arctic Institute of North America, Technical Report, No. 25.
- Hofmann, W. (1971a). Die postglaziale Entwicklung der Chironomiden und *Chaoborus*-Fauna (Dipt.) des Schöhsees. Archiv für Hydrobiologie Supplement 40: 1-74.
- Hofmann, W. (1971b). Zur Taxonomie und Palökologie subfossiler Chironomiden (Dipt.) in Seesedimenten. Archiv für Hydrobiologie Supplement 6: 1-50.
- Hofmann, W. (1978). Analysis of animal microfossils from the Großer Segeberger See (F.R.G.). Archiv für Hydrobiologie 82: 316-346.
- Hofmann, W. (1979). Studies on animal microfossils of sediment cores from northern Germany lakes. International Project on Paleolimnology and Late Cenozoic Climate Newsletter 2: 26.
- Hofmann, W. (1980). Tierische Mikrofossilien aus Oberflächensedimenten einiger Eifelmaare. Mitteilungen der Pollichia 68: 177-184.

- Hofmann, W. (1983a). Stratigraphy of Cladocera and Chironomidae in a core from a shallow North German lake. Hydrobiologia 103: 235-239.
- Hofmann, W. (1983b). Stratigraphy of subfossil Chironomidae and Ceratopogonidae (Insecta: Diptera) in late glacial littoral sediments from Lobsigensee (Swiss Plateau). Studies in the late Quaternary of Lobsigensee 4. Revue de Paléobiologie 2: 205-209.
- Hofmann, W. (1985). Developmental history of Lobsigensee: subfossil Chironomidae (Diptera). pp. 154-156, In: Lobsigensee - Late-glacial and Holocene environments of a lake on the central Swiss plateau. Dissertationes Botanicae 87: 127-170.
- Hofmann, W. (1986). Chironomid analysis. pp. 715-727, In: Handbook of Holocene Palaeoecology and Palaeohydrology. (B.E. Berglund, Ed.), John Wiley & Sons, N.Y.
- Iovino, A.J. (1975). Extant chironomid larval populations and the representativeness and nature of their remains in lake sediments. Ph.D. thesis, Indiana University, Indiana, U.S.A.
- Kangasniemi, B.J., and Oliver, D.R. (1983). Chironomidae (Diptera) associated with *Myriophyllum spicatum* in Okanagan Valley lakes, British Columbia. Canadian Entomologist 115: 1545-1546.
- Kavanaugh, D.H. (1984). Studies on Nebriini (Coleoptera: Carabidae), V. New Nearctic *Nebria* taxa and changes in nomenclature. Proceedings of the California Academy of Sciences 43: 159-177.
- Klinka, K. (1976). Ecosystem units, their classification, interpretation and mapping in the University of British Columbia Research Forest. Ph.D. thesis, Univ. of British Columbia, Vancouver.
- Koskinen, R. (1968). Seasonal and diel emergence of *Chironomus salinarius* Kieff. (Dipt., Chironomidae) near Bergen, western Norway. Annales Zoologici Fennici 5: 65-70.
- Krajina, V.J. (1969). Ecology of forest trees in British Columbia. Ecology of western North America 2(1): 1-146.
- Lawrenz, R.W. (1975). The developmental paleoecology of Green Lake, Antrim County, Michigan. M.S. thesis, Central Michigan University, Michigan, U.S.A.
- Lee, R.L., Jr., and Baust, J.G. (1982). Absence of metabolic cold adaptation and compensatory acclimation in the Antarctic fly, *Belgica antarctica*. Journal of Insect Physiology 28: 725-729.
- Likens, G.E., and Bormann, F.H. (1974). Linkages between terrestrial and aquatic ecosystems. BioScience 24: 447-456.
- Lindeberg, B. (1970). Tanytarsini (Diptera, Chironomidae) from northern Fennoscandia. Annales Zoologici Fennici 7: 303-312.

- Livingstone, D.A. (personal communication). Department of Zoology, Duke University, Durham, North Carolina, U.S.
- Livingstone, D.A. (1953). On the paleolimnology of arctic Alaska. Ph.D. thesis, Yale University, New Haven, Connecticut
- Livingstone, D.A. (1963). Alaska, Yukon, Northwest Territories, and Greenland. pp. 559-574, In: Limnology in North America. (D.G. Frey, Ed.), University of Wisconsin Press, Madison.
- Livingstone, D.A., Bryan, K., Jr., and Leahy, R.G. (1958). Effects of an arctic environment on the origin and development of freshwater lakes. Limnology and Oceanography 3: 192-214.
- Lundbeck, J. (1926). Die Bodentierwelt norddeutscher Seen. Archiv für Hydrobiologie und Planktonkunde, Supplement 7: 1-473.
- MacDonald, G.M., (personal communication). Department of Geography, McMaster University, Hamilton, Ontario.
- MacDonald, G.M., and Ritchie, J.C. (1986). Modern pollen spectra from the western interior of Canada and the interpretation of late Quaternary vegetation development. New Phytologist 103: 245-268.
- MacLean, S.F., Jr. (1975). Ecological adaptations of tundra invertebrates. pp. 269-300, In: Physiological adaptations to the environment. (F.J. Vernberg, Ed.), Intext Educational Publ., New York.
- Maguire, B., Jr. (1963). The passive dispersal of small aquatic organisms and their colonization of isolated bodies of water. Ecological Monographs 33: 161-185.
- Mason, P.G. (1983). Systematics and ecology of Chironomidae (Diptera) associated with Tobin Lake Reservoir and the Saskatchewan River. Ph.D. Thesis, Univ. Saskatchewan.
- Mason, P.G., and Lehmkuhl, D.M. (1983). Effects of the Squaw Rapids Hydroelectric Development on Saskatchewan River Chironomidae (Diptera). Memoirs of the American Entomological Society 34: 187-210.
- Mathewes, R.W. (1973). A palynological study of postglacial vegetation changes in the University Research Forest, southwestern British Columbia. Canadian Journal of Botany 51: 2085-2103.
- Mathewes, R.W. (1985). Paleobotanical evidence for climatic change in southern British Columbia during late-glacial and Holocene time. Syllogeus 55: 397-422.
- Mathewes, R.W. (personal communication). Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia.
- Mathewes, R.W., and Heusser, L.E. (1981). A 12,000 year palynological record of temperature and precipitation trends in southwestern British Columbia. Canadian Journal of Botany 59: 707-710.

- Mathewes, R.W., and Rouse, G.E. (1975). Palynology and paleoecology of postglacial sediments from the lower Fraser River Canyon of British Columbia. Canadian Journal of Earth Sciences 12: 745-756.
- Mathewes, R.W., Vogel, J.S., Southon, J.R., and Nelson, D.E. (1985). Accelerator radiocarbon date confirms early deglaciation of the Queen Charlotte Islands. Canadian Journal of Earth Sciences 22: 790-791.
- Mayhood, D.W., and Anderson, R.S. (1976). Limnological survey of the Lake Louise area, Banff National Park, Part 2 - The lakes. Canadian Wildlife Service, Environment Canada, Calgary.
- Megard, R.O. (1964). Biostratigraphic history of Dead Man Lake, Chuska Mountains, New Mexico. Ecology 45: 529-546.
- Michailova, P., and Fischer, J. (1986). *Chironomus vancouveri* sp.n. from Canada (Diptera, Chironomidae). Reichenbachia 23: 99-106.
- Minns, C.K. (1977). Limnology of some lakes on Truelove Lowland. pp. 569-585, In: Truelove Lowland, Devon Island, Canada: A high arctic ecosystem. (L.C. Bliss, Ed.). University of Alberta Press, Edmonton.
- Moodie, G.E.E., and Reimchen, T.E. (1976). Glacial refugia, endemism, and stickleback populations of the Queen Charlotte Islands, British Columbia. Canadian Field Naturalist 90: 471-474.
- Moore, J.W. (1978). Some factors influencing the diversity and species composition of benthic invertebrate communities in twenty arctic and subarctic lakes. Internationale Revue der gesamten Hydrobiologie 63: 757-771.
- Moore, J.W. (1980). Composition of benthic invertebrate communities in relation to phytoplankton populations in five subarctic lakes. Internationale Revue der gesamten Hydrobiologie 65: 657-671.
- Morley, R.L., and Ring, R.A., (1972a). The intertidal Chironomidae (Diptera) of British Columbia. I. Keys to their life stages. Canadian Entomologist 104: 1093-1098.
- Morley, R.L., and Ring, R.A., (1972b). The intertidal Chironomidae (Diptera) of British Columbia. II. Life history and population dynamics. Canadian Entomologist 104: 1099-1121.
- Mossberg, P., and Nyberg, P. (1979). Bottom fauna of small acid forest lakes. Report of the Institute of Freshwater Research, Drottningholm 58: 77-87.
- Mothes, G. (1968). Einige ökologisch interessante Chironomiden aus dem Stechlinseegebiet. Annales Zoologici Fennici 5: 92-96.
- Murray, D.A. (1983). An inventory of Irish Chironomidae (Diptera). Memoirs of the American Entomological Society 34: 223-233.
- Naumann, E. (1917). Undersökningar öfver fytoplankton och under den pelagiska regionen försiggående gyttjeoch dybildningar inom vissa syo-och mellasvenska urbergsvatten. Kungliga svenska Vetenskapsakademien Handlingar 56: 1-165.

- Naumann, E. (1918). Über die natürliche Nahrung des limnischen Zooplanktons. Lunds universitets årsskrift, ny följd. Andra avdelningen 2. 14: 1-47.
- Naumann, E. (1919). Några synpunkter angående planktons ökologi med särskild hänsyn till fytoplankton. Svensk Botanisk Tidskrift, 13: 129-163. (English translation - Freshwater Biological Assoc. No. 49).
- Naumann, E. (1920). Några synpunkter angående de limniska avlejringsarnes terminologi. Sveriges geologiska Undersökning, Series C. No. 300.
- Northcote, T.G., and Larkin, P.A. (1963). Western Canada. pp. 451-485, In: Limnology in North America. (D.G. Frey, Ed.), University of Wisconsin Press, Madison.
- Nyquist, D., and LaPerriere, J.D. (1973). A preliminary survey of the zooplankton and benthos of an arctic lake near Prudhoe Bay, Alaska. Entomological News 84: 227-234.
- Oliver, D.R. (1963). Entomological studies in the Lake Hazen area, Ellesmere Island, including lists of species of Arachnida, Collembola, and Insecta. Arctic 16: 175-180.
- Oliver, D.R. (1964). A limnological investigation of a large arctic lake, Nettilling Lake, Baffin Island. Arctic 17: 69-83.
- Oliver, D.R. (1968). Adaptations of arctic Chironomidae. Annales Zoologici Fennici 5: 111-118.
- Oliver, D.R. (1971). Life history of the Chironomidae. Annual Review of Entomology 16: 211-230.
- Oliver, D.R. (1976). Chironomidae (Diptera) of Char Lake, Cornwallis Island, N.W.T., with descriptions of two new species. Canadian Entomologist 108: 1053-1064.
- Oliver, D.R. (1981a). Chironomidae. pp. 423-458, In: Manual of Nearctic Diptera, Vol. I., Agriculture Canada Monograph No. 27.
- Oliver, D.R. (1981b). Description of *Euryhapsis* new genus including three new species (Diptera: Chironomidae). Canadian Entomologist 113: 711-722.
- Oliver, D.R. (1983). The larvae of Diamesinae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. Entomologica Scandinavica Supplement 19: 115-138.
- Oliver, D.R. (personal communication). Biosystematics Research Centre, K.W. Neatby Building, Central Experimental Farm, Agriculture Canada, Ottawa, Ontario, Canada K1A 0C6.
- Oliver, D.R., and Danks, H.V. (1972). Sex ratios of some high arctic Chironomidae (Diptera). Canadian Entomologist 104: 1413-1417.
- Oliver, D.R., McClymont, D., and Roussel, M.E. (1978). A key to some larvae of Chironomidae (Diptera) from the Mackenzie and Porcupine River watersheds. Canada Fisheries and Marine Service Technical Report 791: 1-73.

- Oliver, D.R., and Roussel, M.E. (1982). The larvae of *Pagastia* Oliver (Diptera: Chironomidae) with descriptions of three Nearctic species. Canadian Entomologist 114: 849-854.
- Oliver, D.R., and Roussel, M.E. (1983a). The insects and arachnids of Canada. Part 11. The genera of larval midges of Canada; Diptera: Chironomidae. Agriculture Canada Publication 1746: 1-263.
- Oliver, D.R., and Roussel, M.E. (1983b). Redescription of *Brillia* Kieffer (Diptera: Chironomidae) with descriptions of Nearctic species. Canadian Entomologist 115: 257-279.
- Pankratova, V.Y. (1970). Lichinki i kukolki komarov podsemeistva Orthoclaadiinae fauny SSSR (Diptera, Chironomidae=Tendipedidae). (Larvae and pupae of midges of the subfamily Orthoclaadiinae (Diptera, Chironomidae=Tendipedidae) of the USSR fauna). Opred Faune SSSR 102: 1-343.
- Parkinson, A., and Ring, R.A. (1983). Osmoregulation and respiration in a marine chironomid larva, *Paraclunio alaskensis* Coquillett (Diptera, Chironomidae). Canadian Journal of Zoology 61: 1937-1943.
- Paterson, C.G., and Walker, K.F. (1974). Recent history of *Tanytarsus barbatarsus* Freeman (Diptera: Chironomidae) in the sediments of a shallow, saline lake. Australian Journal of Marine and Freshwater Research 25: 315-325.
- Pennak, R.W. (1968). Field and experimental winter limnology of three Colorado mountain lakes. Ecology 49: 505-520.
- Peterson, E.B. (1969). Radiosonde data for characterization of a mountain environment in British Columbia. Ecology 50: 200-205.
- Pinder, L.C.V., and Reiss, F. (1986). The pupae of Chironominae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. Entomologica scandinavica Supplement 28: 299-456.
- Pinder, L.C.V., and Reiss, F. (1983). The larvae of Chironominae (Diptera: Chironomidae) of the Holarctic region - Keys and diagnoses. Entomologica scandinavica Supplement 19: 293-435.
- Pojar, J. (1980). Brooks Peninsula: Possible Pleistocene glacial refugium on northwestern Vancouver Island. Programs and Abstracts, Botanical Society of America, Miscellaneous Publication 158: 89.
- Province of British Columbia. (1967). Forest cover map 103-F-10-d,11-a. Forest Inventory Division, British Columbia Forest Service.
- Province of British Columbia. (1980). Climate of British Columbia, 1977. Air Studies Branch, Assessment and Planning Division, British Columbia Ministry of Environment, Victoria.
- Province of British Columbia. (undated). British Columbia Geological Highway Map. British Columbia Ministry of Energy, Mines, and Petroleum Resources, Victoria.

- Rawson, D.S. (1942). A comparison of some large alpine lakes in western Canada. Ecology 23: 143-161.
- Rawson, D.S. (1955). Morphometry as a dominant factor in the productivity of large lakes. Verhandlungen der internationalen Vereinigung für theoretische und angewandte Limnologie 12: 164-175.
- Reiss, F. (1968). Verbreitung lakustrischer Chironomiden (Diptera) des Alpengebietes. Annales Zoologici Fennici 5: 119-125.
- Ricker, W.E. (1952). The benthos of Cultus Lake. Journal of the Fisheries Research Board of Canada 9: 204-212.
- Rigler, F.H. (1978). Limnology in the high arctic: a case study of Char Lake. Verhandlungen der internationalen Vereinigung für theoretische und angewandte Limnologie 20: 127-140.
- Roback, S.S. (1971). The adults of the subfamily Tanypodinae (=Pelopiinae) in North America (Diptera: Chironomidae). Academy of Natural Sciences of Philadelphia Monograph 17: 1-410.
- Roback, S.S. (1984). Tanypodinae (Diptera: Chironomidae) from Afognak and Kodiak Islands, Alaska. Proceedings, Academy of Natural Sciences, Philadelphia 136: 12-23.
- Roddick, J.A. (1965). Vancouver North, Coquitlam, and Pitt Lake map-areas, British Columbia with special emphasis on the evolution of the plutonic rocks. Geological Survey of Canada Memoir 335: 1-276.
- Rodhe, W. (1969). Crystallization of eutrophication concepts in northern Europe. pp. 50-64, In: Eutrophication: causes, consequences, correctives. Nat. Acad. Sci., Washington, D.C.
- Rossaro, B. (1984). *Stilocladius* Rossaro, 1979 reconsidered, with descriptions of the female and larva of *S. montanus* Rossaro (Diptera: Chironomidae, Orthoclaadiinae). Entomologica scandinavica 15: 185-191.
- Rutter, N.W. (1972). Geomorphology and multiple glaciation in the area of Banff, Alberta. Geological Survey of Canada Bulletin 206: 1-54.
- Sæther, O.A. (1969). Some Nearctic Podonominae, Diamesinae, and Orthoclaadiinae (Diptera: Chironomidae). Bulletin of the Fisheries Research Board of Canada 170: 1-154.
- Sæther, O.A. (1970). A survey of the bottom fauna in lakes of the Okanagan Valley, British Columbia. Fisheries Research Board of Canada Technical Report 196: 1-29.
- Sæther, O.A. (1975a). Nearctic chironomids as indicators of lake typology. Verhandlungen der Internationalen Vereinigung für theoretische und angewandte Limnologie 19: 3127-3133.
- Sæther, O.A. (1975b). Nearctic and Palaearctic *Heterotrissocladius* (Diptera: Chironomidae). Fisheries Research Board of Canada Bulletin 193: 1-67.

- Sæther, O.A. (1975c). Two new species of *Protanypus* Kieffer, with keys to Nearctic and Palearctic species of the genus (Diptera: Chironomidae). Journal of the Fisheries Research Board of Canada 32: 367-388.
- Sæther, O.A. (1975d). Two new species of *Heterotanytarsus* Spärck, with keys to Nearctic and Palearctic males and pupae of the genus (Diptera: Chironomidae). Journal of the Fisheries Research Board of Canada 32: 259-270.
- Sæther, O.A. (1976). Revision of *Hydrobaenus*, *Trissocladius*, *Zalutschia*, *Paratrissocladius* and some related genera (Diptera: Chironomidae). Bulletin of the Fisheries Research Board of Canada 195: 1-287.
- Sæther, O.A. (1977). Taxonomic studies on Chironomidae: *Nanocladius*, *Pseudochironomus*, and the *Harnischia* complex. Fisheries Research Board of Canada Bulletin 196: 1-143.
- Sæther, O.A. (1979). Chironomid communities as water quality indicators. Holarctic Ecology 2: 65-74.
- Sæther, O.A. (1980a). The influence of eutrophication on deep lake benthic invertebrate communities. Progress in Water Technology 12: 161-180.
- Sæther, O.A. (1980b). New name for *Oliveria* Sæther, 1976 (Diptera: Chironomidae) nec *Oliveria* Sutherland, 1965 (†Cnidaria: Anthozoa), with a first record for the European continent. Entomologica scandinavica 11: 399-400.
- Sæther, O.A. (1980c). Glossary of chironomid morphology terminology (Diptera: Chironomidae). Entomologica scandinavica Supplement 14: 1-51.
- Sæther, O.A. (1982). Orthoclaadiinae (Diptera: Chironomidae) from SE U.S.A., with descriptions of *Pludsonia*, *Unniella*, and *Platysmittia* n.gen. and *Atelopodella* n.subgen. Entomologica scandinavica Supplement 13: 465-510.
- Sæther, O.A., and McLean, M.P. (1972). A survey of the bottom fauna in Wood, Kalamalka and Skaha Lakes in the Okanagan Valley, British Columbia. Fisheries Research Board of Canada Technical Report 342: 1-28.
- Sæther, O.A., and Sublette, J.E. (1983). A review of the genera *Doithrix* n.gen., *Georthocladus* Strenzke, *Parachaetocladus* Wülker and *Pseudorthocladus* Goetghebuer (Diptera: Chironomidae, Orthoclaadiinae). Entomologica scandinavica Supplement 20: 1-100.
- Schakau, B., and Frank, C. (1984). Die Entwicklung der Chironomiden-Fauna (Diptera) des Tegeler Sees im Spät- und Postglazial. Verhandlungen der Gesellschaft für Ökologie 12: 375-382.
- Schuster, R.M., and Schofield, W.B. (1982). On *Dendrobazzania*, a new genus of Lepidoziaceae (Jungermanniales). The Bryologist 85: 231-238.
- Simpson, K.W., and Bode, R.W. (1980). Common larvae of Chironomidae (Diptera) from New York state streams and rivers with particular reference to the fauna of artificial substrates. New York State Museum Bulletin 439: 1-105.

- Soponis, A.R. (1977). A revision of the Nearctic species of *Orthocladius* (*Orthocladius*) van der Wulp (Diptera: Chironomidae). Memoirs of the Entomological Society of Canada 102: 1-187.
- Stahl, J.B. (1959). The developmental history of the chironomid and *Chaoborus* faunas of Myers Lake. Investigations of Indiana Lakes and Streams 5: 47-102.
- Stahl, J.B. (1969). The uses of chironomids and other midges in interpreting lake histories. Mitteilungen Internationale Vereinigung für Theoretische und Angewandte Limnologie 17: 111-125.
- Stark, D.M. (1976). Paleolimnology of Elk Lake, Itasca State Park, Northwestern Minnesota. Archiv für Hydrobiologie, Supplement 50: 208-274.
- Sublette, J.E., and Sublette, M.S. (1965). Family Chironomidae (Tendipedidae). pp. 142-181. In: A catalog of the Diptera of America north of Mexico. (A. Stone, C.W. Sabrosky, W.W. Wirth, R.H. Foote, and J.R. Coulson, Eds.), U.S. Department of Agriculture Handbook No. 276, 1696 pp.
- Sutherland Brown, A. (1968). Geology of the Queen Charlotte Islands, British Columbia. British Columbia Department of Mines and Petroleum Resources, Bulletin No. 54.
- Thienemann, A. (1915). Die Chironomidenfauna der Eifelmaare. Verhandlungen des Naturhistorischer Verein der Rheinlande und Westfalens 72: 1-58.
- Thienemann, A. (1918). Untersuchungen über die Beziehungen zwischen dem Sauerstoffgehalt des Wassers und der Zusammensetzung der Fauna in Norddeutschen Seen. Archiv für Hydrobiologie 12: 1-65.
- Thienemann, A. (1921). Seetypen. Die Naturwissenschaften 18: 343-346.
- Thomasson, K. (1956). Reflections on arctic and alpine lakes. Oikos 7: 117-143.
- Thut, R.N. (1969). A study of the profundal bottom fauna of Lake Washington. Ecological Monographs 39: 79-100.
- Timms, B.V., Hammer, U.T., and Sheard, J.W. (1986). A study of benthic communities of some saline lakes in Saskatchewan and Alberta, Canada. Internationalen Revue der gesamten Hydrobiologie 71: 759-777.
- Tobolski, K. (1985). Plant macrofossils from Lobsigensee. pp. 140-143. In: Lobsigensee - Late-glacial and Holocene environments of a lake on the central Swiss plateau. Dissertationes Botanicae 87: 127-170.
- Topping, M.S. (1969). Giant chromosomes, ecology, and adaptation in *Chironomus tentans*. Ph.D. thesis, University of British Columbia, Vancouver.
- Topping, M.S. (1971). Ecology of larvae of *Chironomus tentans* (Diptera: Chironomidae) in saline lakes in central British Columbia. Canadian Entomologist 103: 328-338.

- Tsukada, M. (1982). *Pseudotsuga menziesii* (Mirb.) Franco: its pollen dispersal and late Quaternary history in the Pacific Northwest. Japanese Journal of Ecology 32: 159-187.
- Tuiskunen, J., and Lindeberg, B. (1986). Chironomidae (Diptera) from Fennoscandia north of 68° N, with a description of ten new species and two new genera. Annales zoologici Fennici 23: 361-393.
- Vitt, D.H., and Schofield, W.B. (1976). *Seligeria careyana*, a new species from the Queen Charlotte Islands, western Canada. The Bryologist 79: 231-234.
- Wainman, N. (personal communication). Surrey Campus, Kwantlen College, Surrey, British Columbia
- Wainman, N., and Mathewes, R.W. (1987). Forest history of the last 12 000 years based on plant macrofossil analysis of sediment from Marion Lake, southwestern British Columbia. Canadian Journal of Botany 65: 2179-2187.
- Walker, I.R. (1987). Chironomidae (Diptera) in paleoecology. Quaternary Science Reviews 6: 29-40.
- Walker, I.R., Fernando, C.H., and Paterson, C.G. (1984). The chironomid fauna of four shallow humic lakes and their representation by subfossil assemblages in the surficial sediments. Hydrobiologia 112: 61-67.
- Walker, I.R., Fernando, C.H., and Paterson, C.G. (1985). Associations of Chironomidae (Diptera) of shallow, acid, humic lakes and bog pools in Atlantic Canada, and a comparison with an earlier paleoecological investigation. Hydrobiologia 120: 11-22.
- Walker, I.R., and Mathewes, R.W. (1987a). Chironomidae (Diptera) and postglacial climate at Marion Lake, British Columbia, Canada. Quaternary Research 27: 89-102.
- Walker, I.R., and Mathewes, R.W. (1987b). Chironomids, lake trophic status, and climate. Quaternary Research 28: 431-437.
- Walker, I.R., and Paterson, C.G. (1983). Post-glacial chironomid succession in two small, humic lakes in the New Brunswick - Nova Scotia (Canada) border area. Freshwater Invertebrate Biology 2: 61-73.
- Warner, B.G. (1984). Late Quaternary paleoecology of eastern Graham Island, Queen Charlotte Islands, British Columbia, Canada. Ph.D. thesis, Simon Fraser University, Burnaby, British Columbia.
- Warner, B.G., and Chmielewski, J.G. (1987). Biometric analysis of modern and late Pleistocene cones of *Picea* from western Canada. New Phytologist 107: 449-457.
- Warner, B.G., Clague, J.J., and Mathewes, R.W. (1984). Geology and paleoecology of a mid-Wisconsin peat from the Queen Charlotte Islands, British Columbia, Canada. Quaternary Research 21: 337-350.

- Warner, B.G., Mathewes, R.W., and Clague, J.J. (1982). Ice-free conditions on the Queen Charlotte Islands, British Columbia, at the height of late Wisconsin glaciation. Science 218: 675-677.
- Warwick, W.F. (1975). The impact of man on the Bay of Quinte, Lake Ontario, as shown by the subfossil chironomid succession (Chironomidae, Diptera). Verhandlungen der internationalen Vereinigung für theoretische und angewandte Limnologie 19: 3134-3141.
- Warwick, W.F. (1980a). Chironomidae (Diptera) responses to 2800 years of cultural influence; a palaeolimnological study with special reference to sedimentation, eutrophication, and contamination processes. Canadian Entomologist 112: 1193-1238.
- Warwick, W.F. (1980b). Palaeolimnology of the Bay of Quinte, Lake Ontario: 2800 years of cultural influence. Canadian Bulletin of Fisheries and Aquatic Sciences 206: 1-117.
- Warwick, W.F. (personal communication). Environment Canada, National Hydrological Laboratory, 11 Innovation Blvd, Saskatoon, Saskatchewan S7N 2X8
- Weber, C.A. (1907). Aufbau und Vegetation der Moore Norddeutschlands. Beiblatt zu den Botanischen Jahrbuchern 90: 19-34.
- Welch, H.E., Jr. (1973). Emergence of Chironomidae (Diptera) from Char Lake, Resolute, Northwest Territories. Canadian Journal of Zoology 51: 1113-1123.
- Welch, H.E., Jr. (1976). Ecology of Chironomidae (Diptera) in a polar lake. Journal of the Fisheries Research Board of Canada 33: 227-247.
- Wetzel, R.G. (1975). Limnology. W.B. Saunders, Philadelphia 743 pp.
- Whiteside, M.C. (1983). The mythical concept of eutrophication. Hydrobiologia 103: 107-111.
- Wiederholm, T. (1976). A survey of the bottom fauna of Lake Sammamish. Northwest Science 50: 23-31.
- Wiederholm, T. (1979). Chironomid remains in recent sediments of Lake Washington. Northwest Science 53: 251-256.
- Wiederholm, T. (1980). Effects of dilution on the benthos of an alkaline lake. Hydrobiologia 68: 199-207.
- Wiederholm, T. (Ed.). (1983). Chironomidae of the Holarctic region. Keys and diagnoses. Part 1 - Larvae. Entomologica Scandinavica Supplement 19: 1-457.
- Wiens, A.P., Rosenberg, D.M., and Snow, N.B. (1975). Species list of aquatic plants and animals collected from the Mackenzie and Porcupine River watersheds from 1971-1973. Canada Fisheries and Marine Service Technical Report 557: 1-39.

- Winnell, M.H., and White, D.S. (1985). Ecology of some Chironomidae (Diptera) from southeastern Lake Michigan, U.S.A. Transactions of the American Entomological Society 111: 279-359.
- Wirth, W.W., and Sublette, J.E. (1970). A review of the Podonominae of North America with descriptions of three new species of *Trichotanypus* (Diptera: Chironomidae). Journal of the Kansas Entomological Society 43: 335-354.
- Zetterstedt, J.W. (1838). Sectio tertia. Diptera Dipterologis scandinaviae amices et popularibus carissimus. pp. 477-868, In: *Insecta lapponica*. Leipzig.